

# Metric Spaces Worksheet 5

## Topology I

With our understanding of metric spaces and sequences cemented, we'll turn to examine a notion which is supported by every metric space, and in some ways subsumes the concepts we have seen so far.

**Definition 1** (open ball). Let  $(X, d)$  be a metric space,  $x \in X$  a point and  $r \in (0, \infty)$  a non-negative real number. The *open ball of radius  $r$  centred on  $x$* , written  $B_r(x)$ , is the subset  $B_r(x) := \{y \in X \mid d(x, y) < r\} \subseteq X$ .  $\square$

We now calculate open balls in Euclidean metric spaces. To describe open balls in the Euclidean line, we need the notion of an *open interval* in  $\mathbb{R}$ . For any  $a, b \in \mathbb{R}$ , with  $a < b$ , let

$$(a, b) := \{z \in \mathbb{R} \mid a < z < b\}.$$

### Example 2 (open balls in Euclidean spaces)

1. In the Euclidean metric space  $\mathbb{R}$ , the open ball  $B_r(x) = \{y \in \mathbb{R} \mid |x - y| < r\}$  is the open interval  $(x - r, x + r)$ . Conversely, every open interval  $(a, b)$  for  $a < b \in \mathbb{R}$ , is an open ball of some radius  $r = \frac{b-a}{2}$  centred about the midpoint  $\frac{a+b}{2}$ .
2. In the Euclidean metric space  $\mathbb{R}^2$ , the open ball

$$B_r((u_1, u_2)) = \{(v_1, v_2) \in \mathbb{R}^2 \mid (u_1 - v_1)^2 + (u_2 - v_2)^2 < r^2\}$$

is comprised of all points inside the circle of radius  $r$  centred at the point  $(u_1, u_2)$ . This explains the name “open ball” given to the sets  $B_r(x)$  in general metric spaces.

**Question 3.** What are the possible open balls in a discrete metric space  $(X, d)$ ?

Complete the proof here



**Definition 4** (open set). A subset  $U \subseteq X$  in a metric space  $(X, d)$  is *open* if for every  $u \in U$  there exists an  $\varepsilon \in (0, \infty)$  such that  $B_\varepsilon(u) \subseteq U$ .  $\lrcorner$

**Example 5** (*an open set in the Euclidean space  $\mathbb{R}$* )

For any  $a \in \mathbb{R}$ , the open ray

$$(a, \infty) := \{x \in \mathbb{R} \mid a < x\}$$

is an open set.

To see this we must prove, for every point  $u \in (a, \infty)$ , that there exists some  $\varepsilon \in (0, \infty)$  such that  $B_\varepsilon(u) \subseteq (a, \infty)$ . To that end, consider a point  $u \in (a, \infty)$ . We know that  $u - a > 0$ , so we may choose  $\varepsilon$  to be any real number so that  $0 < \varepsilon < u - a$ . (For sake of concreteness, we might pick  $\varepsilon = \frac{u-a}{2}$ , but it's also not necessary to specify a concrete value of  $\varepsilon$ .)

Now if  $x \in B_\varepsilon(u)$ , then by example 2 item 1,  $u - \varepsilon < x < u + \varepsilon$ . Since  $u - \varepsilon > a$  we conclude that  $x > a$  so  $x \in (a, \infty)$ . Since we've shown that  $\forall x \in B_\varepsilon(u), x \in (a, \infty)$  this demonstrates that  $B_\varepsilon(u) \subseteq (a, \infty)$  as required. Thus  $(a, \infty)$  is an open set.

**Non-example 6** (*sets which are not open in the Euclidean space  $\mathbb{R}$* )

1. The set  $\{0\} \subseteq \mathbb{R}$  is not open because there is no  $\varepsilon$  small enough so that  $B_\varepsilon(0) \subset \{0\}$ .
2. For any  $a \in \mathbb{R}$ , the closed ray

$$[a, \infty) := \{x \in \mathbb{R} \mid a \leq x\}$$

is not an open set. The argument given in example 5 proves that for every  $u \in [a, \infty)$  if  $a \neq u$  then there exists  $\varepsilon \in (0, \infty)$  so that  $B_\varepsilon(u) \subset [a, \infty)$ . However, there is no open ball that contains the point  $a$  and is contained within  $[a, \infty)$ .

To see this, take  $\varepsilon \in (0, \infty)$ . Then by example 2 item 1 the point  $a - \frac{\varepsilon}{2} \in B_\varepsilon(a)$ . But since  $a - \frac{\varepsilon}{2} < a$ ,  $a - \frac{\varepsilon}{2} \notin [a, \infty)$ . Thus  $B_\varepsilon(a) \not\subseteq [a, \infty)$ .

**Question 7.** What are the open sets in a discrete metric space  $(X, d)$ ?


Complete the proof here



As we might hope, the subsets we were calling *open* balls are indeed open.

**Proposition 8** (open balls are open sets). *Let  $(X, d)$  be a metric space,  $x \in X$  be a point,  $r \in (0, \infty)$  be a non-negative real number. The subset  $B_r(x) \subseteq X$  is open.*

Complete the proof here



**Corollary 9** (open intervals are open sets). *In the Euclidean metric space  $\mathbb{R}$ , all open intervals  $(a, b)$  are open.*

Complete the proof here



It turns out that open sets can be combined in certain ways and the result is always again an open set.

**Theorem 10** (open set laws). *In a metric space  $(X, d)$ ,*

1.  $X$  and  $\emptyset$  are open sets.
2. If  $\mathcal{F}$  is a family of open sets in  $X$  then  $\cup_{U \in \mathcal{F}} U$  is open.
3. If  $U, V \subseteq X$  are open sets then  $U \cap V$  is open.

Complete the proof here

**Surprise 11 (*intersection of opens is not generally open*)**

In the Euclidean metric space  $\mathbb{R}$ , the subset  $I := \bigcap_{n \in \mathbb{N}} \left(0, \frac{n+2}{n+1}\right) \subseteq \mathbb{R}$  is not open.

Compute  $I$  and prove this fact.

Complete the proof here



Created by tslil clingman, 2019. This work is licensed under a Creative Commons “Attribution-ShareAlike 4.0 International” license.

